

## REVIEW

# Breathing retraining and exercise conditioning in patients with chronic obstructive pulmonary disease (COPD): a physiological approach

F. GIGLIOTTI, I. ROMAGNOLI AND G. SCANO

Fondazione Don C. Gnocchi (IRCCS), Pozzolatico, Firenze, Italy

**Abstract** In this review we shall consider the commonest techniques to reduce dyspnea that are being applied to patients with chronic obstructive pulmonary disease (COPD) subjected to a pulmonary rehabilitation program (PRP). Pursed lip breathing (PLB) and diaphragmatic breathing (DB) are breathing retraining strategies employed by COPD patients in order to relieve and control dyspnea. However, the effectiveness of PLB in reducing dyspnoea is controversial. Moreover, DB may be associated with asynchronous and paradoxical breathing movements, reflecting a decrease in the efficiency of the diaphragm. Exercise training (EXT) is a mandatory component of PRP. EXT has been shown to improve exercise performances and peripheral muscle strength. Recent studies have focused on the effect of EXT on breathlessness. However, concerns persist as to whether the decreased sensation of dyspnea for a given exercise stimulus is principally due to psychological benefits of rehabilitation or to improved physiological ability to perform exercise. The effect of EXT on breathlessness may be reinforced by inhaling oxygen. However, two studies have recently shown that breathing supplemental oxygen during training has either a marginal effect or no advantage over training. In a comprehensive PRP, strength training (ST) and arm endurance training (AET) could have a role in decreasing peripheral muscle weakness and metabolic and ventilatory requirements for AET. The role of unloading the respiratory muscles during EXT has to be clarified. © 2002 Elsevier Science Ltd. All rights reserved.

Available online at <http://www.sciencedirect.com>

**Keywords** pulmonary rehabilitation; dyspnea; COPD.

## INTRODUCTION

One of the two major goals in the management of chronic obstructive pulmonary disease (COPD) is to decrease respiratory symptoms and improve the quality of life (QoL). As a result of pulmonary rehabilitation programs (PRP), improvements have been demonstrated in objective measures of QoL (1), well-being (2) and health status, including reduction of respiratory symptoms, increase in exercise tolerance and functional activities such as walking, enhanced ability of performing activities of daily living and improved psychological function (3,4).

Referral to a comprehensive pulmonary rehabilitation program is indicated in COPD patients who have been placed on optimal medical therapy and who: (1) continue

to display severe respiratory symptoms, mostly breathlessness; (2) have had several emergency ward or hospital admissions per year; (3) exhibit limited functional status and restricted activities of daily living and (4) experience impairment in QoL (3,4).

PRPs are usually employed on an outpatient basis and comprehensive programs vary in their frequency and duration, with the optimal frequency and length remaining a matter of debate.

In this review, we shall consider the most common techniques to reduce dyspnea that are being applied to patients with COPD.

## BREATHING RETRAINING (BR) AND BODY POSTURE

The goal of BR is to help the patient to relieve and control breathlessness and counteract abnormalities such as dynamic hyperinflation associated with COPD.

Received 24 April 2002, accepted in revised form 8 August 2002  
Correspondence should be addressed to: Giorgio Scano, Fondazione Don C. Gnocchi (IRCCS), Pozzolatico, Via Imprunetana 124, 50020 Firenze, Italy. E-mail: [g.scano@dmf.unifi.it](mailto:g.scano@dmf.unifi.it)

Review of clinical studies evaluating the effects of BR indicate that improvement in clinical symptoms is a more consistent finding than any measurable impact on physiological parameters (3,4).

### Pursed lip breathing (PLB)

PLB is a breathing retraining strategy often spontaneously and voluntarily employed by COPD patients in order to relieve and control dyspnea during exercise or daily activities or during periods of increased ventilatory demand (5,6). PLB improves gas exchange (5–8), decreases respiratory rate, increases tidal volume (5,7,9,10) and increases the activity of inspiratory and expiratory muscles that take over the activity of breathing (5). In circumstances of hyperinflation, an increased motor command to, and a reduced muscle strength of, the diaphragm are likely explanations for dyspnea (11). A deflationary activity on chest wall of PLB should potentially reduce breathlessness. However, the effectiveness of PLB in reducing dyspnea in COPD is controversial, with some studies demonstrating either decrease (5,8,12), no changes (13) or increase in breathlessness at rest (10) and during exercise (14). The fact that PLB does not promote any reduction in pulmonary volume and whether the self-imposed rhythmic respiration with PLB affects chest wall (CW) motion and compartmental coordinated activity have as yet to be demonstrated. By applying a 3-D optoelectronic plethysmograph (OEP), Nerini *et al.* (15) have recently shown a marked decrease in end expiratory lung volume (EELV), localized at abdominal level, a pattern never shown before. They have also confirmed previous data by Breslin (6) showing increase in expiratory muscle activity, and have extended those data in that abdominal muscle activity is greater than that derived from the mere measures of changes in end expiratory gastric pressure ( $P_{gae}$ ).

Anecdotal evidence of impaired breathlessness with PLB has been reported in COPD patients (5,14). We speculate that this may in part be due to increased activity of the respiratory muscles aimed at avoiding overtaxing the diaphragm.

### Leaning forward (LF)

Although this is not a part of PRP, it may help patients to relieve breathlessness. Standing position increases EMG of the diaphragm ( $E_{di}$ ), lowers trans-diaphragmatic pressure ( $P_{di}$ ) and increases  $E_{di}/P_{di}$  ratio. At variance, LF decreases EMG activity of many respiratory muscles (16), including the diaphragm, and improves  $E_{di}/P_{di}$  ratio (17). The reason for the association of LF with decrease in breathlessness lies in the common belief that breathlessness is linked to the increase in the central motor command to the respiratory muscles (18).

Decrease in respiratory muscle activation lowers breathlessness.

### Diaphragmatic breathing (DB)

The costal and crural parts of the muscle are electrically arranged partially in series and partially in parallel (19). With hyperinflation, costal and crural parts are arranged in parallel, a condition where: (1) the zone of apposition decreases; (2) the abdominal pressure ( $P_{ab}$ ) no longer expands the rib cage (RC); (3) the system produces less pressure; (4) the diaphragm expands the abdomen but has a deflationary action on the RC. Thus, in patients with COPD DB may be associated with asynchronous and paradoxical breathing movements (20,21). A recent study by Gosselink *et al.* (22) showed that, compared to quiet breathing, with DB inspiratory abdominal volume ( $V_{ab}$ ) increases while the volume of rib cage ( $V_{rc}$ ) decreases. This is a paradoxical breathing reflecting the limitation of DB which decreases the efficiency of the diaphragm. In the same study, dyspnea tended to increase with DB. Recently similar results have been produced by Vitacca *et al.* (23) in hypercapnic patients with COPD. It is possible that patients with limited hyperinflation may benefit from DB while severely hyperinflated patients are incapable of performing DB (24).

Association of breathing retraining with body position is being commonly performed by patients to reduce dyspnea (3,4).


### EXERCISE TRAINING (EXT)

EXT is a mandatory component of PRP. Patients with COPD should regularly perform aerobic lower extremity endurance exercises to enhance performance of daily activities and reduce dyspnea (3,4).

#### General exercise training

Patients experience increased capacity and endurance for exercise and physical activity after EXT even though lung function may remain unchanged (25–28). Recent studies have shown an early onset of lactic acidosis during exercise at high exercise levels in COPD patients (29–32). Improvements in maximal and submaximal exercise responses obtained after EXT at high exercise levels (30,32,33) indicate improved aerobic metabolism (30,32,33). The study by Casaburi *et al.* (30) showing the effects of sustained level of exercise on the delay in anaerobic threshold in COPD patients with moderate airway obstruction indicates that a physiological training response had occurred. Maltais *et al.* (32) have indicated a beneficial effect on skeletal muscle level as shown by the increase in lactate threshold, citrate synthase (CS) and 3-hydroxy-acyl CoA dehydrogenase (HADH). Reduction

Casaburi *et al.* (41) have focused on the mechanism whereby EXT improves exercise tolerance in COPD (Fig. 1). On the other hand recent studies have focused on the effect of EXT on breathlessness. The disparity between the respiratory motor output and the mechanical response of the system is thought to play a major role in the increase perception of exercise dyspnea in COPD patients (49,50). Conflicting results, however, indicate the lack of relationship between dyspnea and any mechanical variable in exercising COPD (51). Therefore, concerns



More recently, O'Donnell *et al.* (47) have shown that general EXT improves strength and endurance of the peripheral leg muscles, decreases  $B_{org}$  and log effort (LE) but strength and endurance do not relate to either  $B_{org}$  or LE. So increased tolerance to stimuli or altered perceptual response to evoked sensation may also contribute to score symptom alleviation. Nonetheless, evidence of increased symptom tolerance associated with EXT should be provided after close evaluation of the mechanical factors involved in the coupling between respiratory motor output and mechanical response of the respiratory system (55).

The effect of EXT on breathlessness may be reinforced by inhaling oxygen. Administration of 30–60% oxygen

enhances ventilatory exercise performance in COPD (56–60). Casaburi *et al.* (30) have shown that decrease in VE was associated with decrease in lactate production. Inhaling 60% oxygen reduced lactate production and ventilation in COPD (60). As a consequence of the improved aerobic metabolism, breathlessness decreases at iso-work load (58). Decrease in VE is the reason for decrease in breathlessness, particularly in patients with more severe degree of obstruction and hypoxemia (60). Former studies by Criner and Celli (61) with 30% oxygen administration had shown a change in the strategy of respiratory muscle recruitment whereby increase in diaphragmatic performance avoids overtaxing RC and accessory muscles and decreases breathlessness. Nevertheless, the effect of oxygen on breathlessness may occur independently from ventilatory changes indicating the role of central mechanism(s) on the perception of breathlessness (62). Two recent studies (63,64) have recently shown that breathing supplemental oxygen during training has either a marginal effect or no advantage over training while breathing room-air. Despite the evidence of short-term beneficial effects, there are no studies on long-term effects of oxygen on factors limiting effort tolerance and breathlessness in patients with COPD.

## TRAINING TO STRENGTH AND ENDURANCE OF THE INSPIRATORY MUSCLES

Inspiratory muscle training (IMT) in addition to exercise training has been shown to improve exercise capacity more than exercise training alone (65–67). IMT as monotherapy was found to decrease dyspnea (67–69). Strength training may be obtained by maximum static inspiratory (MIP) and expiratory effort over vital capacity every 3–5 min for periods of 20–30 min a day. Alternatively, strength can be also increased by loading protocol involving inspiratory effort against flow resistive or threshold loads to generate target levels of peak inspiratory mouth pressure  $> 33\%$  MIP for 30–60 min daily for 5–7 days a week (69,70). The lack of improvement in MIP noted in some trials may be because of insufficient training intensity (71). Indeed, a recent meta-analysis has shown that overall exercise endurance improved in studies of ventilatory muscle training in which the training stimulus led to an increase in MIP but did not improve in the studies without improvements in respiratory muscle strength following training (72).

Endurance and the sense of respiratory effort depend on pressure, inspiratory flow, inspiratory time ( $T_i$ ), and the ratio of  $T_i$  to the total time of the respiratory cycle ( $T_{tot}$ ), and  $R_f$  (73). It seems probable that subjects breathing spontaneously on a given resistance may vary their inspiratory flow rate decreasing the sense of respiratory

effort by reducing peak inspiratory pressure ( $P_i$ ), or the pressure time index ( $PT_i$ ), i.e., the area under the pressure–time curve of the inspiratory muscles. In this way, peak intra-thoracic pressure and  $PT_i$  may be similar with and without resistance. In other words, resistive breathing has the disadvantage that inspiratory pressure is flow-dependent. For training to have consistent effects on endurance, patients have to be coached to breathe with longer  $T_i$  and lower peak pressure such that  $PT_i$  is greater than during spontaneous breathing (74). In that study, while peak mouth pressure was higher during spontaneous breathing,  $PT_i$  was appreciably greater during coached breathing. After a training period of 10 weeks maintaining  $PT_i$  at 25%,  $P_i/P_{i\max}$  at 50% and  $T_i/T_{tot}$  at 50%, Criner *et al.* (75) found an increase in trans-diaphragmatic ( $P_{di}$ ) twitch of about 40%. Breathing on resistance loads at high respiratory rate increases strength and endurance in a small number of COPD patients.

Unlike resistive breathing, isocapnic hyperventilation (70–90% maximum voluntary ventilation (MVV)) for periods of 15-min a day 6 times per week (76,77) is associated with lower levels of pressure but greater inspiratory flow and probably a greater extent of muscle shortening. Thus, isocapnic hyperventilation increases velocity of shortening at low level of pressure or tension (78). Like isocapnic hyperventilation, threshold load (TL) is independent of flow rate. Also, TL enhances the velocity of inspiratory muscle contraction, shortening inspiratory time and increasing time for exhalation and relaxation. However, at high loads decrease in  $T_i$  for a given  $P_{di}$  reduces  $T_i/T_{tot}$  thereby reducing  $P_{di}$ , which may be the most important training variable. Despite improvement in respiratory muscle strength and endurance (79,80), with relief in breathlessness in some patients (67–69), IMT may not result in enhanced exercise performance (79).

## TRAINING OF PERIPHERAL MUSCLES

### Strength training (ST)

ST is obtained by additional weights to lower and upper limb movement. The importance of peripheral skeletal muscle dysfunction in the impairment of exercise capacity and LE in patients with COPD was suggested by Killian and colleagues (81,82). Training of peripheral muscle strength has been shown to improve maximal muscle strength, exercise endurance capacity and QoL but not maximal exercise capacity (83,84). In healthy subjects, strength training may (85) or may not (86) enhance the effect of endurance training. In patients with COPD, the addition of strength training to endurance training has no additional effect on exercise performance and QoL (87). However, in patients with muscle weakness the combination of strength with endurance appears to

**TABLE I.** Summary of scientific evidence grades for the described techniques

Program contents	Recommendations	Grade
Lower extremity training	Lower extremity training improves exercise tolerance and is recommended as a part of pulmonary rehabilitation	A
Upper extremity training	Strength and endurance training improves arm function; arm exercises should be included in pulmonary rehabilitation	B
Ventilatory muscle training	Is not an essential component of pulmonary rehabilitation	B
Oxygen administration	Does not enhance rehabilitation outcomes	D
Breathing retraining	No advantages.	D
Respiratory muscle rest	Can enhance skeletal muscle training	C

Modified from refs. (70, 102).

increase peripheral muscle strength, exercise performance and QoL (48).

### Endurance arm exercise training (AET)

AET involves a larger muscle mass working at moderate intensity for a longer period of time. For equal work rates, ventilation and oxygen consumption are generally higher for arm than for leg exercise (88,89). This is probably due to a lower mechanic efficiency of the arm muscle because more static work is needed to stabilize trunk and shoulder. It likely follows an earlier onset of anaerobic metabolism for the arms than for the legs. The effect of unsupported arm exercise (UAET) on respiratory system increases by increasing arm loading (90). In severely obstructed patients, AET is associated with dyssynchronous breathing and is terminated because of severe dyspnea, this not being observed with the same patients performing more intense leg (cycle) exercise, with a higher  $\dot{V}_{O_2}$ , heart rate and exercise duration (91). The reason for dyssynchronous breathing with AET was thought to be due to a decreased contribution of RC and accessory muscles to ventilation overtaxing the diaphragm and leading to its earlier fatigue. Several lines of evidence indicate that unsupported AET increases  $\dot{V}_{O_2}$ ,  $\dot{V}_{CO_2}$  output ( $\dot{V}_{CO_2}$ ), end inspiratory gastric pressure ( $P_{gai}$ ) and both end expiratory pleural ( $P_{ese}$ ) and gastric ( $P_{gae}$ ) pressures in patients with CAO (92). Compared to UAET, supported AET (SAET) decreases  $P_{gai}$  and  $P_{esi}$  pressures and  $P_{gae}$  in most of the COPD patients again indicating that a greater part of VE during UAET is shifted from RC inspiratory muscles to both the diaphragm and the expiratory muscles.

Comprehensive pulmonary rehabilitation including arm exercise decreases metabolic and ventilatory requirement for AET (93). The reduction of  $\dot{V}_{O_2}$  during arm activity which results from such training might be expected to improve dyspnea (1). Upper extremity (UE) training program improves specific UE performances (38,93–95) but not activities simulating daily living (38,94) or walking test (95).

### UNLOADING THE RESPIRATORY MUSCLES

Maltais *et al.* (96) showed that 11 cmH<sub>2</sub>O of inspiratory pressure support reduces both inspiratory effort and dyspnea in severe CAO patients performing constant work load bicycle exercise. In particular, the inverse relationship between improvement in dyspnea and decrease in pressure–time integral of both inspiratory muscles and the diaphragm, i.e., decrease in inspiratory effort was evident. Furthermore, there was no further change in EELV with addition of PS compared with control exercise. The authors maintained that, unlike continuous positive airway pressure (CPAP), the efficacy of inspiratory support do not necessarily depend on either the prevailing level of intrinsic positive end expiratory pressure (PEEP<sub>i</sub>) or the presence of expiratory FL.

O'Donnell *et al.* (97) were able to show that CPAP decreased dyspnea in exercising patients with CAO. They postulated that unloading the respiratory muscles resulted in a decreased central respiratory output and thereby a decreased inspiratory effort, in as much as, in exercising patients trans-pulmonary pressure promotes airway compression thus contributing to dyspnea. The application of 4 cmH<sub>2</sub>O positive end expiratory pressure (PEEP) contributed to dyspnea relief by attenuating dynamic compression (97). In two recent papers, Harms *et al.* (98,99) have shown that unloading the respiratory muscles with proportional assist ventilation during strenuous exercise in cyclists reduces oxygen uptake and the perception of both breathlessness and leg discomfort, indicating that work of breathing significantly influences exercise performance. The effect of the normal respiratory muscle load on exercise performance in trained cyclists may be due to the associated reduction in leg blood flow which increases both leg fatigue and the intensity with which leg effort and respiratory muscle effort are perceived. This also explains the difficulty of discriminating between the two sensations. The link between respiratory work and exercise performance is likely to be due to a vasoconstrictor effect from the diaphragm to the limb muscle vasculature. This

occurs during heavy exercise with sustained work of breathing and cardiac output limited in its ability to distribute flow adequately to both respiratory and locomotor muscles. The same may happen in conditions of moderate exercise when cardiac output is abnormally low, likely to occur in exercising patients with severe COPD. Recently, Garrod *et al.* (100) have shown that the addition of non-invasive pressure positive ventilation (NPPV) to an exercise training program in severe COPD results in a significant improvement in mean shuttle walk test. At variance, Bianchi *et al.* (101) have recently shown that proportional assist ventilation during training sessions included in a multidisciplinary PRP is not well tolerated by all patients and gave no additional physiological benefit in comparison with exercise training alone (Table I).

## REFERENCES

- Guyatt GH, Berman LB, Townshed M, *et al.* A measure of quality of life for clinical trials in chronic lung disease. *Thorax* 1987; **42**: 773–778.
- Atkins CJ, Kaplan RM, Timms RM, *et al.* Behavioral exercise programs in the management of chronic obstructive pulmonary disease. *J Consult Clin Psychol* 1984; **52**: 59–603.
- American Thoracic Society. Standards for the diagnosis and care of patients with chronic obstructive pulmonary disease. ATS statement. *Am Rev Respir Crit Care Med* 1995; **152**(Suppl): S77–S120.
- American Thoracic Society. Pulmonary rehabilitation—1999. *Am J Respir Crit Care Med* 1999; **159**: 1666–1682.
- Mueller RE, Petty TL, Filley GF. Ventilation and arterial blood gas changes induced by pursed lips breathing. *J Appl Physiol* 1970; **28**: 784–789.
- Breslin EH. The pattern of respiratory muscle recruitment during pursed-lip breathing. *Chest* 1992; **101**: 75–78.
- Casciari RJ, Fairshirer RD, Harrison A, *et al.* Effects of breathing retraining in patients with chronic obstructive pulmonary disease. *Chest* 1981; **79**: 393–398.
- Ingram RH, Schilder DP. Effect of pursed-lips expiration on the pulmonary pressure flow relationship in obstructive lung disease. *Am Rev Respir Dis* 1967; **96**: 381–383.
- Thoman RL, Stoker GL, Ross JC. The efficiency of pursed-lips breathing in patients with obstructive pulmonary disease. *Am Rev Respir Dis* 1966; **93**: 100–106.
- Breslin EH, Ugalde V, Bonekat HW, *et al.* Abdominal muscle recruitment during pursed-lip breathing in COPD (abstract). *Am J Respir Crit Care Med* 1996; **153**: A128.
- Killian KJ, Gandevia SC, Summers WV, *et al.* Effect of increased lung volume on perception of breathlessness effort and tension. *J Appl Physiol* 1984; **57**: 686–691.
- Sharp J, Danon J, Druz W, *et al.* Respiratory muscle function in patients with chronic obstructive pulmonary disease: its relationship to disability and to respiratory therapy. *Am Rev Respir Dis* 1974; **110**: 154–167.
- Roa J, Epstein S, Breslin E, Celli B. Work of breathing and ventilatory muscle recruitment during pursed lips breathing. *Am Rev Respir Dis* 1991; **143**: A77.
- Spahija J, de Marchie M, Grassino A. Pursed-lips breathing during exercise increases dyspnea (abstract). *Am Rev Respir Dis* 1993; **147**: A729.
- Nerini M, Gigliotti F, Bianchi R, *et al.* Changes in global and compartmental lung volumes during pursed lip breathing (PLB) in COPD patients. *Eur Respir J* 2001; **18** (Suppl. 33): 489s (Abstract).
- Sharp JT, Druz WS, Moisan T, *et al.* Postural relief of dyspnea in severe chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1980; **122**: 201–211.
- Druz WS, Sharp JT. Electrical and mechanical activity of the diaphragm accompanying body position in severe chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1982; **125**: 275–280.
- Killian KJ, Jones NL. Respiratory muscles and dyspnea. *Clin Chest Med* 1988; **9**: 237–248.
- Macklem PT, Macklem DM, De Troyer A. A model of inspiratory muscle mechanics. *Respir Environ Exerc Physiol* 1983; **55**(2): 547–557.
- Willeput R, Vachaud J P, Lenders D, *et al.* Thoraco-abdominal motion during chest physiotherapy in patients affected by chronic obstructive lung disease. *Respiration* 1983; **44**: 204–214.
- Sackner MA, Silva G, Banks JM, *et al.* Distribution of ventilation during diaphragmatic breathing in obstructive lung disease. *Am Rev Respir Dis* 1974; **109**: 331–337.
- Gosselink R, Wagenaar RC, Sargeant AJ, *et al.* Diaphragmatic breathing reduces efficiency of breathing in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1995; **151**: 1136–1142.
- Vitacca M, Clini E, Bianchi L, Ambrosino N. Acute effects of deep diaphragmatic breathing in COPD patients with chronic respiratory insufficiency. *Eur Respir J* 1998; **11**: 408–415.
- Breslin EH. Breathing retraining in COPD. *J Cardiopulm Rehabil* 1995; **15**: 25–33.
- Lertzman MM, Cherniack RM. Rehabilitation of patients with chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1976; **114**: 1145–1165.
- Toshima MT, Ries AL. Experimental evaluation of rehabilitation in chronic obstructive pulmonary disease: short term effects on exercise endurance and health status. *Health Psychol* 1990; **9**: 237–252.
- McGavin CR, Gupta SP, McHardy GJR. Twelve-minute walking test for assessing disability in chronic bronchitis. *Br Med J* 1976; **1**: 822–823.
- Casaburi R. Exercise training in chronic obstructive lung disease. In: Casaburi R, Petty TL (eds). *Principles and Practice of Pulmonary Rehabilitation*. Philadelphia, PA: Saunders, 1993: 204–224.
- Casaburi R, Wasserman K, Patessio A, *et al.* A new perspective in pulmonary rehabilitation: anaerobic threshold as a discriminant in training. *Eur Respir J* 1989; **2**: 618s–623s.
- Casaburi R, Patessio A, Ioli F, *et al.* Reduction in exercise lactic acidosis and ventilation as a result of exercise training in obstructive lung disease. *Am Rev Respir Dis* 1991; **143**: 9–18.
- Patessio A, Casaburi R, Carone M, *et al.* Comparison of gas exchange, lactate, and lactic acidosis threshold in patients with chronic obstructive pulmonary disease. *Am rev Respir Dis* 1993; **148**: 622–626.
- Maltais F, Leblanc P, Simard C, *et al.* Skeletal muscle adaptation to endurance training in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1996; **154**: 442–447.
- Maltais F, Leblanc P, Jobin J, *et al.* Intensity of training and physiologic adaptation in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1997; **155**: 555–561.
- Foster S, Lopez D, Thomas HM III. Pulmonary rehabilitation in COPD patients with elevated PCO<sub>2</sub>. *Am Rev Respir Dis* 1988; **138**: 1519–1523.
- Niederman M, Clemente P, Fein A, *et al.* Benefits of a multi-disciplinary pulmonary rehabilitation program: improvements are independent of lung function. *Chest* 1991; **99**: 798–804.

36. Davis JA, Whipp BJ, Lamarra N. Effect of ramp slope on determination of aerobic parameters from the ramp exercise test. *Med Sci Sports Exerc* 1982; **14**: 335–343.
37. Nery LE, Wasserman K, Andrews JD, et al. Ventilatory and gas exchange kinetics during exercise in chronic airway obstruction. *J Appl Physiol* 1982; **53**: 1594–1602.
38. Ries AL, Kaplan RM, Limberg TM, et al. Effects of pulmonary rehabilitation on physiologic and psychosocial outcomes in patients with chronic obstructive pulmonary disease. *Ann Intern Med* 1995; **122**: 823–832.
39. Ries AL and the ACCP/AACVPR Pulmonary Rehabilitation Guidelines Panel. Pulmonary Rehabilitation: Joint ACCP/ AACVPR evidence-based guidelines. *J Cardiopulm Rehabil* 1997; **17**: 371–405.
40. Simpson K, Killian KJ, McCartney N, et al. Randomized controlled trial of weightlifting exercise in patients with chronic airflow limitation. *Thorax* 1992; **47**: 70–75.
41. Casaburi R, J Porszasz, Burns MR, et al. Physiologic benefits of exercise training in rehabilitation of patients with severe chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1997; **155**: 1541–1551.
42. O'Donnell DE, McGuire MA, Samis L, Webb KA. The impact of exercise reconditioning on breathlessness in severe chronic airflow limitation. *Am J Respir Crit Care Med* 1995; **152**: 2005–2013.
43. Goldstein RS, Gort EH, Stubbing D, et al. Randomized controlled trial of respiratory rehabilitation. *Lancet* 1994; **344**: 1394–1397.
44. Casaburi R. Special considerations for exercise training in chronic lung disease. In: *ACSM's Resource Manual for Guidelines for Exercise Training and Prescription*, 3rd edn. Williams & Wilkins, Baltimore, MD, 1998: 334–338.
45. American College of Sports Medicine. Position stand: the recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness in healthy adults. *Med Sci Sports Exerc* 1990; **22**: 265–274.
46. Gosselink R, Troosters T, Decramer M. Exercise training in COPD: interval training versus endurance training. *Eur Respir J* 1998; **12** (Suppl. 28): 2s.
47. O'Donnell DE, McGuire MA, Samis IL, Webb KA. General exercise training improves ventilatory and peripheral muscle strength and endurance in chronic airflow limitation. *Am J Respir Crit Care Med* 1998; **157**: 1489–1497.
48. Gosselink R, Troosters T, Decramer M. Physiological adaptation after exercise training in patients with COPD. *Am J Respir Crit Care Med* 1997; **155**: A 497.
49. O'Donnell DE. Breathlessness in patients with chronic airflow limitations. Mechanisms and management. *Chest* 1994; **106**: 904–912.
50. Belman MJ, Botnick WVC, Shin JW. Inhaled bronchodilators reduce dynamic hyperinflation during exercise in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1996; **153**: 967–975.
51. Montes de Oca M, Rassulo J, Celli BR. Respiratory muscle and cardiopulmonary function during exercise in very severe COPD. *Am J Respir Crit Care Med* 1996; **154**: 1284–1289.
52. Belman MJ, Kendregan BA. Exercise training fails to increase skeletal muscle enzymes in patients with chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1981; **123**: 256–261.
53. Haas F, Salazar-Schicchi, Axen K. Desensitization to dyspnea in chronic obstructive pulmonary disease. In: Casaburi R, Petty TL (eds) *Principles and Practice of Pulmonary Rehabilitation*. Philadelphia, PA: Saunders 1993: 241–251.
54. Sinclair JD. The effect of breathing exercise in pulmonary emphysema. *Thorax* 1955; **10**: 246–249.
55. Rosi E, Gigliotti F, Misuri G, et al. Effects of exercise training (ET) on exertional dyspnoea in severe COPD patients. *Eur Respir J* 2000; **16**: P370.
56. Scano G, Van Meerhaeghe A, Willeput R et al. Effect of oxygen on breathing during exercise in patients with chronic lung disease. *Eur J Respir Dis* 1982; **63**: 23–30.
57. Somfay A, Porszasz J, Lee SM, Casaburi R. Dose–response effect of oxygen on hyperinflation and exercise endurance in nonhypoxaemic COPD patients. *Eur Respir J* 2001; **18**: 77–84.
58. Dean NC, Brown JK, Himelman RB, et al. Oxygen may improve dyspnea and endurance in patients with chronic obstructive pulmonary disease and only hypoxemia. *Am Rev Respir Dis* 1992; **148**: 941–945.
59. Bye PTP, Esau SA, Levy RD, et al. Ventilatory muscle function during exercise in air and oxygen in patients with chronic airflow limitation. *Am Rev Respir Dis* 1985; **132**: 236–240.
60. O'Donnell DE, D'Arsigny C, Webb K. Effects of hyperoxia on ventilatory limitation during exercise in advanced chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2001; **163**: 892–898.
61. Criner GJ, Celli BR. Ventilatory muscle recruitment in exercise with O<sub>2</sub> in obstructed patients with mild hypoxemia. *J Appl Physiol* 1987; **63**: 195–200.
62. Lane R, Cockcroft A, Adams L, Guz A. Arterial oxygen saturation and breathlessness in patients with chronic obstructive airway disease. *Clin Sci* 1987; **72**: 693–698.
63. Garrod R, Paul EA, Wedzicha JA. Supplemental oxygen during pulmonary rehabilitation in patients with COPD with exercise hypoxemia. *Thorax* 2000; **55**: 539–543.
64. Rooyackers JM, Dekhuizen PNR, Van Herwaarden CLA, Folgering HTM. Training with supplemental oxygen in patients with COPD and hypoxemia at peak exercise. *Eur Respir J* 1997; **10**: 1278–1284.
65. Dekhuijzen PNR, Folgering HTM, van Herwaarden CLA. Target-flow inspiratory muscle training during pulmonary rehabilitation in patients with COPD. *Chest* 1991; **99**: 128–133.
66. Wanke TH, Formanek D, Lahrman H, et al. Effects of combined inspiratory muscle and cycle ergometer training on exercise performance in patients with COPD. *Eur Respir J* 1994; **7**: 2205–2211.
67. Lisboa C, Villafranca C, Leiva A, et al. Inspiratory muscle training in chronic airflow limitation: effect on exercise performance. *Eur Respir J* 1997; **10**: 537–542.
68. Lisboa C, Munoz V, Beroiza T, et al. Inspiratory muscle training in chronic airflow limitation: comparison of two different training loads with a threshold device. *Eur Respir J* 1994; **7**: 1266–1274.
69. Harver A, Mahler DA, Daubenspeck JA. Targeted inspiratory muscle training improves respiratory muscle function and reduces dyspnea in patients with chronic obstructive pulmonary disease. *Ann Intern Med* 1989; **111**: 117–124.
70. American College of Chest Physicians and American Association for Cardiovascular and Pulmonary Rehabilitation. Pulmonary Rehabilitation Joint ACCP and AACVPR evidence-based guidelines. *Chest* 1997; **112**: 1363–1396.
71. Preusser BA, Winningham MI, Clanton TL. High versus low intensity inspiratory muscle interval training in patients with COPD. *Chest* 1994; **106**: 110–117.
72. Smith K, Cook D, Guyatt G, et al. Respiratory muscle training in chronic airflow limitation: a meta-analysis. *Am Rev Respir Dis* 1992; **145**: 533–539.
73. Jones GL, Killian KJ, Summers E, Jones NL. Inspiratory muscles forces and endurance in maximum resistive loading. *J Appl Physiol* 1985; **58**: 1608–1621.
74. Belman MJ, Thomas SF, Lewis MI. Resistive breathing training in patients with chronic obstructive pulmonary disease. *Chest* 1986; **90**: 663–670.
75. Criner GJ, Travaline JM, Sudarshan S, Leyenson V. Effect of inspiratory muscle training on diaphragm strength and fatigability. *Am J Respir Crit Care Med* 1995; **151**: A807.

76. Leith DE, Bradley M. Ventilatory muscle strength and endurance training. *J Appl Physiol* 1976; **41**: 508–516.
77. Belman MJ, Mittman C. Ventilatory muscle training improves exercise capacity in chronic obstructive pulmonary disease patients. *Am Rev Respir Dis* 1980; **121**: 273–280.
78. Duchateau J, Hainaut K. Isometric or dynamic training: differential effects on mechanical properties of a human muscle. *J Appl Physiol* 1984; **56**: 296–301.
79. Larson JL, Covey MK, Wirtz SE, *et al.* Cycle ergometry and inspiratory muscle training in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1999; **160**: 500–507.
80. Villafranca C, Borzone G, Leiva A, *et al.* Effect of inspiratory muscle training with an intermediate load on inspiratory power output in COPD. *Eur Respir J* 1998; **11**: 28–33.
81. Killian KJ, LeBlanc P, Martin DH, *et al.* Exercise capacity and ventilatory, circulatory, symptom limitation in patients with chronic airflow limitation. *Am Rev Respir Dis* 1992; **146**: 935–940.
82. Hamilton N, Killian KJ, Summers E, Jones NL. Muscle strength, symptom intensity, and exercise capacity in patients with cardiorespiratory disorders. *Am J Respir Crit Care Med* 1995; **152**: 2021–2031.
83. Simpson K, Killian K, McCartney N, *et al.* Randomized controlled trial of weightlifting exercise in patients with chronic airflow limitation. *Thorax* 1992; **47**: 70–75.
84. Clark CJ, Cochrane JE, Mackay E. Low intensity peripheral muscle conditioning improves exercise tolerance and breathlessness in COPD. *Eur Respir J* 1996; **9**: 2590–2596.
85. Hickson RC, Dvorak BA, Gorostiaga EM, *et al.* Potential for strength and endurance training to amplify endurance performance. *J Appl Physiol* 1988; **65**: 2285–2290.
86. Dudley GA, Djamil R. Incompatibility of endurance- and strength-training modes of exercise. *J Appl Physiol* 1985; **59**: 1446–1451.
87. Bernard S, Whittom F, Leblanc P, *et al.* Aerobic and strength training in patients with chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1999; **159**: 897–901.
88. Toner MM, Sawka MN, Levine L, Pandolf KB. Cardiorespiratory responses to exercise distributed between upper and lower body. *J Appl Physiol* 1983; **54**: 1403–1407.
89. Casaburi R, Barstow TJ, Robinson T, Wasserman K. Dynamic and steady-state ventilatory and gas exchange responses to arm exercise. *Med Sci Sports Exerc* 1992; **24**: 1365–74.
90. Romagnoli I, Gorini M, Gigliotti F, *et al.* Chest-wall kinematics and respiratory muscle recruitment during unsupported arm exercise in healthy humans. Submitted to World Congress on Lung Health and 12th ERS Annual Congress 2002. *Eur Respir J* 2002; **20**: 3015.
91. Celli BR, Rassulo J, Make BJ. Dyssynchronous breathing during arm but not leg exercise in patients with chronic airflow obstruction. *N Engl J Med* 1986; **314**: 1485–1490.
92. Martinez FJ, Couser JI, Celli BR. Respiratory response to arm elevation in patients with chronic airflow obstruction. *Am Rev Respir Dis* 1991; **143**: 476–480.
93. Couser JI, Martinez FJ, Celli BR. Pulmonary rehabilitation that includes arm exercise reduced metabolic and ventilatory requirements for simple arm elevation. *Chest* 1993; **103**: 37–41.
94. Lake FR, Henderson K, Briffa T, *et al.* Upper-limb and lower-limb exercise training in patients with chronic airflow obstruction. *Chest* 1990; **97**: 1077–1082.
95. Gosselink R, Decramer M. Muscle training in pulmonary rehabilitation. *Eur Respir Monogr Pulm Rehabil* 2000; **5**: 99–110.
96. Maltais F, Reissman H, Gottfried SB. Pressure support reduces inspiratory effort and dyspnea during exercise in chronic airflow obstruction. *Am J Respir Crit Care Med* 1995; **151**: 1027–1033.
97. O'Donnell DE, Sanij R, Giesbrecht G, Younes M. Effect of continuous positive airway pressure on respiratory sensation in patients with chronic obstructive pulmonary disease during sub maximal exercise. *Am Rev Respir Dis* 1988; **138**: 1185–1191.
98. Harms CA, Babcock MA, McClaran SR, *et al.* Respiratory muscle work compromises leg blood flow during maximal exercise. *J Appl Physiol* 1997; **82**: 1573–1583.
99. Harms CA, Wetter TJ, St Croix CM, *et al.* Effects of respiratory muscle work on exercise performance. *J Appl Physiol* 2000; **89**: 131–138.
100. Garrod R, Mikelsons C, Paul EA, Wedzicha JA. Randomised controlled trial of domiciliary noninvasive positive pressure ventilation and physical training in severe chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2000; **162**: 1335–1341.
101. Bianchi L, Foglio K, Porta R, *et al.* Lack of additional effect of adjunct of assist ventilation to pulmonary rehabilitation in mild COPD patients. *Respir Med* 2002; **96**: 359–367.
102. British Thoracic Society Standards of Care Subcommittee on Pulmonary Rehabilitation. (Pulmonary rehabilitation.) *Thorax* 2001; **56**: 827–834.